

- 43 Verrillo RT, Bolanowski SJ, Checkoski CM, *et al.* Effects of hydration on tactile sensation. *Somatosens Mot Res* 1998;**15**:93–108.
- 44 Vega- Bermudez F, Johnson KO. Fingertip skin conformance accounts ,in parts, for differences in tactile acuity in young subjects, but not for the decline in spatial acuity with ageing. *Percept Psychophys* 2004;**66**:60–7.
- 45 Downie AW, Newell DJ. Sensory nerve conduction in patients with diabetes mellitus and controls. *Neurology (Minneapolis)* 1961;**11**:876–82.
- 46 Rivner MH, Swift TR, Malik K. Influence of age and height on nerve conduction. *Muscle Nerve* 2001;**24**:1134–41.
- 47 Nadler MA, Harrison LM, Stephens JA. Changes in cutaneomuscular reflex responses in relation to normal ageing in man. *Exp Brain Res* 2002;**146**:48–53.
- 48 Wohler AB. Tactile perception of spatial stimuli on the lip surface by young and older adults. *J Speech Hear Res* 1996;**39**:1191–8.
- 49 Petrosino L, Fucci D, Robey RR. Changes in lingual sensitivity as a function of age and stimulus exposure time. *Percept Mot Skills* 1982;**55**:1083–90.
- 50 Newman HF. Palatal sensitivity to touch: correlation with age. *J Am Geriatr Soc* 1979;**27**:319.
- 51 Canetta R. Decline in oral perception from 20 to 70 years. *Percept Mot Skills* 1977;**45**:1028–30.
- 52 Dannenbaum RM, Jones LA. The assessment and treatment of patients who have sensory loss following cortical lesions. *J Hand Ther* 1993;**6**:130–8.
- 53 Tremblay F, Wong, Sanderson R, *et al.* Tactile spatial acuity in elderly persons: assessment with grating domes and relationship with manual dexterity. *Somatosens Mot Res* 2003;**20**:127–32.
- 54 Ranganathan VK, Siemionow V, Sahgal V, *et al.* Effects of ageing on hand function. *J Am Geriatr Soc* 2001;**49**:1478–84.
- 55 Lord SR, Clark RD, Webster IW. Postural stability and associated physiological factors in a population of aged persons. *J Gerontol* 1991;**46**:69–76.
- 56 Tanaka T, Noriyasu S, Ino S, *et al.* Objective method to determine the contribution of the great toe to standing balance and preliminary observations of age-related effects. *IEEE Trans Rehabil Eng* 1996;**4**:84–90.
- 57 Corriveau H, Hebert R, Raiche M, *et al.* Postural stability in the elderly: empirical confirmation of a theoretical model. *Arch Gerontol Geriatr* 2004;**39**:163–77.
- 58 Tremblay F, Mireault AC, Dessureault L, *et al.* Postural stabilization from fingertip contact: I. Variations in sway attenuation , perceived stability and contact forces with ageing. *Exp Brain Res* 2004;**157**:275–85.

Clinical Evidence—Call for contributors

Clinical Evidence is a regularly updated evidence-based journal available worldwide both as a paper version and on the internet. *Clinical Evidence* needs to recruit a number of new contributors. Contributors are healthcare professionals or epidemiologists with experience in evidence-based medicine and the ability to write in a concise and structured way.

Areas for which we are currently seeking contributors:

- Pregnancy and childbirth
- Endocrine disorders
- Palliative care
- Tropical diseases

We are also looking for contributors for existing topics. For full details on what these topics are please visit www.clinicalevidence.com/ceweb/contribute/index.jsp

However, we are always looking for others, so do not let this list discourage you.

Being a contributor involves:

- Selecting from a validated, screened search (performed by in-house Information Specialists) epidemiologically sound studies for inclusion.
 - Documenting your decisions about which studies to include on an inclusion and exclusion form, which we keep on file.
 - Writing the text to a highly structured template (about 1500-3000 words), using evidence from the final studies chosen, within 8-10 weeks of receiving the literature search.
 - Working with *Clinical Evidence* editors to ensure that the final text meets epidemiological and style standards.
 - Updating the text every 12 months using any new, sound evidence that becomes available.
- The *Clinical Evidence* in-house team will conduct the searches for contributors; your task is simply to filter out high quality studies and incorporate them in the existing text.

If you would like to become a contributor for *Clinical Evidence* or require more information about what this involves please send your contact details and a copy of your CV, clearly stating the clinical area you are interested in, to CECommissioning@bmjgroup.com.

Call for peer reviewers

Clinical Evidence also needs to recruit a number of new peer reviewers specifically with an interest in the clinical areas stated above, and also others related to general practice. Peer reviewers are healthcare professionals or epidemiologists with experience in evidence-based medicine. As a peer reviewer you would be asked for your views on the clinical relevance, validity, and accessibility of specific topics within the journal, and their usefulness to the intended audience (international generalists and healthcare professionals, possibly with limited statistical knowledge). Topics are usually 1500-3000 words in length and we would ask you to review between 2-5 topics per year. The peer review process takes place throughout the year, and out turnaround time for each review is ideally 10-14 days.

If you are interested in becoming a peer reviewer for *Clinical Evidence*, please complete the peer review questionnaire at www.clinicalevidence.com/ceweb/contribute/peerreviewer.jsp